

Albumin-Bilirubin Score: A Novel Mortality Predictor in Valvular Surgery

Zihni Mert Duman^{1*}, MD; Barış Timur^{2*}, MD

¹Department of Cardiovascular Surgery, Cizre State Hospital, Şırnak Turkey.

²Department of Cardiovascular Surgery, Dr Siyami Ersek Thoracic and Cardiovascular Surgery Training and Research Hospital, İstanbul, Turkey.

*Both authors were working at Mehmet Akif Ersoy Hospital when the study was conducted.

This study was carried out at the Istanbul Mehmet Akif Ersoy Thoracic and Cardiovascular Surgery Training and Research Hospital, Istanbul, Turkey.

ABSTRACT

Introduction: The heart and liver are two organs that are closely related. The Albumin-Bilirubin (ALBI) score is a developed scoring system for assessing liver function. The aims of this study were to examine the correlation between preoperative ALBI score and pulmonary artery pressure and to investigate its ability to predict heart valve surgery mortality outcomes.

Methods: The data of 872 patients who underwent isolated and combined heart valve surgery from 2014 to 2021 were retrospectively screened. In the preoperative period, 152 patients with laboratory tests including albumin and total bilirubin were found and analyzed retrospectively. Thirteen of these patients were excluded from the study. The remaining 139 patients were included in the analysis. Baseline demographic data, echocardiography data, performance status, laboratory data, operative data, and postoperative status

were collected. The optimal cutoff value of preoperative ALBI score was calculated.

Results: The cutoff for ALBI scores was calculated as -2.44 to predict in-hospital mortality (sensitivity = 75.0%, specificity = 70%). Based on the cutoff value, 90 patients had a low ALBI score (≤ -2.44 , 64.7%) and 49 patients had a high ALBI score (> -2.44 , 35.3%). High ALBI score was associated with an increased incidence of acute kidney injury and in-hospital mortality, and a positive correlation was found between ALBI score and pulmonary artery pressure.

Conclusion: In patients with valvular surgery, high ALBI score was an independent prognostic factor of in-hospital mortality and acute kidney injury. It is easily measurable and a cost-effective way to predict mortality.

Keywords: Heart Valve. Acute Kidney Injury. Albumins. Aptitude. Echocardiography. Hospital Mortality. Preoperative Period.

Abbreviations, Acronyms & Symbols

AF	= Atrial fibrillation	EuroSCORE	= European System for Cardiac Operative Risk Evaluation
AKI	= Acute kidney injury	Hba1c	= Glycosylated hemoglobin
ALBI	= Albumin-Bilirubin	Hgb	= Hemoglobin
ALT	= Alanine aminotransferase	IABP	= Intra-aortic balloon pump
APTT	= Activated partial thromboplastin time	ICU	= Intensive care unit
AST	= Aspartate aminotransferase	INR	= International normalized ratio
AUC	= Area under the ROC curve	LVEF	= Left ventricular ejection fraction
BUN	= Blood urea nitrogen	MELD	= Model for End-Stage Liver Disease
CC	= Cross-clamping	NYHA	= New York Heart Association
CI	= Confidence interval	OR	= Odds ratio
COPD	= Chronic obstructive pulmonary disease	PAP	= Pulmonary artery pressure
CPB	= Cardiopulmonary bypass	PLT	= Platelet
CRE	= Creatinine	ROC	= Receiver operating characteristic
DM	= Diabetes mellitus	WBC	= White blood cell
ECMO	= Extracorporeal membrane oxygenation		

Correspondence Address:

Barış Timur

 <https://orcid.org/0000-0003-4446-6374>

Dr. Siyami Ersek Göğüs Kalp ve Damar Cerrahisi Eğitim ve Araştırma Hastanesi

Tıbbiye cad, No: 13, Üsküdar, İstanbul, Turkey

Zip Code: 34668

E-mail: dr.baristimur@gmail.com

Article received on January 7th, 2022.

Article accepted on May 31st, 2022.

Accepted on March 25th, 2026. Erratum available.

INTRODUCTION

The heart and liver are two organs that are closely related. Liver failure may develop in patients with right heart failure due to pulmonary arterial hypertension^[1]. The clinical and prognostic value of the interaction between the heart and liver is not clearly known, and some studies show that the Model for End-Stage Liver Disease (MELD) score, which is used for risk assessment in patients with liver failure, can also be used for risk assessment in cardiac surgery patients^[2-4].

The Albumin-Bilirubin (ALBI) score was recommended by Johnson et al.^[5] as an alternative to the Child-Pugh (or C-P) grade and the MELD score for risk assessment of liver function and subsequent long-term mortality in patients with liver disease. The ALBI score is a simpler test used to evaluate liver function that includes only serum albumin and bilirubin levels. This score is easier to calculate than the MELD score, and one of its advantages is that it is not affected by warfarin usage. In addition, the ALBI score is closely associated with hospital mortality in patients with heart failure as demonstrated in recent studies^[6,7]. The ALBI score may be appropriate for preoperative risk analysis and evaluation of right heart failure due to pulmonary arterial hypertension in patients undergoing heart valve surgery.

The aims of this study were to examine the correlation between preoperative ALBI score and pulmonary artery pressure (PAP) and to investigate its ability to predict heart valve surgery mortality outcomes.

METHODS

Patients who underwent only valvular surgery were included in the study. The data of 872 patients who had undergone isolated and combined heart valve surgery from 2014 to 2021 in Istanbul Mehmet Akif Ersoy Thoracic and Cardiovascular Surgery Training and Research Hospital were retrospectively screened. In the preoperative period, 152 patients with laboratory tests including albumin and total bilirubin were found and analyzed retrospectively. Exclusion criteria for this study were patients with known liver disease including positive hepatitis B antigen and anti-hepatitis C virus antigen and the patients who had to be operated in shock state or sepsis. Thirteen of these patients were excluded from the study, because they satisfied at least one of the exclusion criteria. The remaining 139 patients were included in the study to be analyzed. The baseline demographic data, echocardiographic data, performance status, laboratory data, operative data, and postoperative status were comprehensively collected and analyzed. PAP values were estimated from echocardiography.

The formula used to calculate the ALBI score is: $(\text{albumin} \times -0.085) + (\log_{10} \text{bilirubin} \times 0.66)$, where albumin is measured in g/L and bilirubin in $\mu\text{mol/L}$, as it was previously described in the literature^[5]. The primary outcome was in-hospital mortality. Secondary outcomes included other postoperative complications, acute kidney injury (AKI), pneumonia, and re-exploration for bleeding. We used Standard Society of Thoracic Surgeons definitions in the study^[8]. Hospital mortality was defined as mortality occurring within 30 days postoperatively or without discharge.

Statistical Analysis

Statistical analyses were carried out using IBM Corp. Released 2015, IBM SPSS Statistics for Windows, version 23.0, Armonk, NY: IBM

Corp. Descriptive statistics are reported as percentage for categorical variables and mean $\pm$ standard deviation for continuous variables. Categorical variables were compared by a chi-squared analysis or Fisher's exact test. Normal and abnormal continuous variables were compared by Student's *t*-test and Mann-Whitney U test. Receiver operating characteristic (ROC) analysis was performed to find the appropriate cutoff value for the preoperative ALBI score. The ability of the ALBI score to predict hospital mortality was assessed using the area under the ROC curve (AUC). Univariate analysis of hospital mortality was performed using logistic regression model. Multivariate analysis was performed with variables that were statistically significant in univariate analysis. The correlation between the variables was analyzed using the Pearson's or Spearman's correlation coefficient. Linear regression model was used for the relationship between PAP and ALBI score. Statistical tests were two-sided, and *P*-values < 0.05 were considered statistically significant.

Our study was found ethically appropriate according to the decision of the Health Sciences University Mehmet Akif Ersoy Training and Research Hospital Clinical Research Ethics Committee (dated 04.06.2021; file number 2021/49).

RESULTS

After randomization and exclusion of patients, the remaining 139 patients were included into the study. Most of the patients were women (56.1%) and mean age was 55.6 ± 12.51 years. Three valves in 14 (10.1%) patients, two valves in 70 (50.4%) patients, and one valve in 55 (39.5%) patients were either replaced or repaired. Fifteen patients undergone cardiac reoperation (10.8%). Table 1 shows different operation types that were performed on these patients.

Mean ALBI score was -2.53 ± 0.57 and median ALBI score was -2.59 . Pearson's correlation coefficients were used to analyze the relationship between ALBI score and PAP. We found a positive correlation ($r: 0.245$, $P=0.004$) between ALBI score and PAP. By linear regression analysis, we found that every 1 increase in ALBI score resulted in an increase in PAP of 7.95 mmHg ($P=0.015$). The graph is shown in Figure 1.



In-hospital mortality occurred in 16 (11.5%) patients. ROC curve analysis was established using the preoperative ALBI score to predict in-hospital mortality. The cutoff for ALBI scores was calculated as -2.44 for predicting in-hospital mortality (sensitivity = 75.0%, specificity = 70%, likelihood ratio: 2.5). The AUC was 0.712 and *P*-value was 0.004 for ROC curve analysis. Figure 2 shows that the preoperative ALBI score had a significant positive relationship with in-hospital mortality.

Seventeen of the 139 patients in the study were operated for infective endocarditis. Among patients with infective endocarditis, in-hospital mortality occurred in four patients. ROC curve analysis was done using the preoperative ALBI score to predict in-hospital mortality, excluding patients with infective endocarditis. The AUC was 0.697 and *P*-value was 0.025 for ROC curve analysis for this group of patients.

According to the cutoff value, 49 patients had a high ALBI score ($\text{ALBI} > -2.44$, 35.3%) and 90 patients had a low ALBI score ($\text{ALBI} \leq -2.44$, 64.7%). Preoperative laboratory values and demographic data of patients with low and high ALBI scores are shown in Table 2. Patients with New York Heart Association (NYHA) class 3-4 symptoms were more common in the high ALBI score group

Table 1. Distribution of operation types.

Type of surgery	N (%)
Isolated aortic valve surgery	15 (10.8%)
Isolated mitral valve surgery	38 (27.3%)
Combined mitral and tricuspid valve surgery	50 (36%)
Combined aortic and mitral valve surgery	20 (14.4%)
Combined aortic, mitral, and tricuspid valve surgery	16 (11.5%)
Reoperation	15 (10.8%)

**Fig. 1** - Correlation between the Albumin-Bilirubin (ALBI) score and the pulmonary artery pressure (PAP).**Fig. 2** - Receiver operating characteristic (ROC) analysis of optimal albumin-bilirubin value.

($P < 0.001$). Patients with a high European System for Cardiac Operative Risk Evaluation (EuroSCORE) II were more common in the group with high ALBI scores ($P < 0.037$). Also, patients who had valve surgery for infective endocarditis were more common in the high ALBI score group ($P = 0.007$). Although there was no difference in ejection fraction between the two groups in echocardiographic data, PAPs were higher in the group with higher ALBI scores ($P = 0.011$). Compared to the preoperative laboratory values, only the international normalized ratio values were higher in the high ALBI score group ($P = 0.048$). The rest of the demographical and laboratory data were statistically insignificant ($P > 0.05$).

Perioperative data of patients with low and high ALBI scores are shown in Table 3. There was no statistical difference between re-exploration for bleeding, cross-clamping time, and cardiopulmonary bypass (CPB) time ($P > 0.05$).

Univariable and multivariable analyses were performed to identify independent risk factors related to in-hospital mortality. High preoperative ALBI score (odds ratio [OR]: 3.83, $P = 0.004$), NYHA class 3-4 symptoms (OR: 4.51, $P = 0.013$), high EuroSCORE II (OR: 1.25, $P = 0.002$), CPB time (OR: 1.02, $P < 0.001$), and combined aortic, mitral, and tricuspid valve surgery (OR: 6.78, $P = 0.002$) were associated with hospital mortality in univariate analysis. Multivariate analysis using the stepwise model shows that high preoperative ALBI score (OR: 3.37, $P = 0.036$) and CPB time (OR: 1.02, $P = 0.001$) are independently associated with hospital mortality. Univariate and multivariate mortality analyses are detailed in Table 4.

Higher ALBI score was associated with an increased incidence of AKI ($P = 0.028$) and hospital mortality ($P < 0.001$). However, no significant differences were found for other complications ($P > 0.05$). Table 5 shows the results of postoperative complications according to ALBI score of -2.44.

DISCUSSION

There are problems in liver functions due to chronic congestive hepatopathy in heart valve patients^[9]. Abnormalities in liver function tests in the preoperative period affect the surgical condition of patients. Preoperative risk analysis and prediction of prognosis in patients are the cornerstones of surgical management. Problems in liver function tests are important risk factors that are not considered in the risk models used in cardiac surgery^[10]. In our study, we used the ALBI score, a newly developed method for risk assessment of patients with liver disease, to evaluate the preoperative risk for mortality in heart valve surgery patients. The ALBI score, which uses only serum albumin and total bilirubin values, can be easily measured^[5]. As demonstrated by other studies on the prognosis of liver diseases, the ALBI score is useful in determining the degree of liver dysfunction^[11,12].

Table 2. Preoperative demographic and clinical characteristics of the patients according to their ALBI scores.

	ALBI score < -2.44 (n=90) N (%) or mean±standard deviation	ALBI score > -2.44 (n=49) N (%) or mean±standard deviation	P-value
Age (years)	55.32±12.665	56.12±12.360	0.805
Sex (female)	54 (60%)	24 (48.9%)	0.211
NYHA class 3-4	36 (40%)	29 (59.1%)	0.00*
Preoperative atrial fibrillation	22 (24.4%)	18 (36.7%)	0.148
Infective endocarditis	6 (6.6%)	11 (22.4%)	0.007*
LVEF	54.77±9.44	55.20±9.62	0.727
PAP	40.91±17.31	48.86±19.66	0.011*
EuroSCORE II	2.40±1.88	4.22±5.11	0.037*
Reoperation	8 (9%)	7 (14%)	0.327
DM	22 (24.4%)	9 (18.3%)	0.411
Renal failure	13 (14.4%)	9 (18.3%)	0.545
COPD	22 (24.4%)	12 (24.5%)	0.995

Preoperative laboratory values

WBC (109/L)	8.08±1.99	9.63±3.48	0.244
Hgb	19.51±11.42	17.10±11.28	0.313
PLT	236.88±63.87	279.08±57.55	0.090
ALT (U/L)	22.33±15.57	30.04±29.96	0.226
AST (U/L)	24.11±13.27	35.33±36.482	0.143
BUN	20.32±8.95	26.46±19.721	0.159
CRE	1.16±1.87	1.20±1.76	0.502
Hba1c	5.95±1.65	5.81±1.31	0.761
APTT (s)	29.88±6.46	30.16±5.71	0.301
INR	1.32±0.74	1.41±0.59	0.048*

*P-value < 0.05 is statistically significant

ALBI=Albumin-Bilirubin; ALT=alanine aminotransferase; APTT=activated partial thromboplastin time; AST=aspartate aminotransferase; BUN=blood urea nitrogen; COPD=chronic obstructive pulmonary disease; CRE=creatinine; DM=diabetes mellitus; EuroSCORE=European System for Cardiac Operative Risk Evaluation; Hba1c=glycosylated hemoglobin; Hgb=hemoglobin; INR=international normalized ratio; LVEF=left ventricular ejection fraction; NYHA=New York Heart Association; PAP=pulmonary artery pressure; PLT=platelet; WBC=white blood cell

Table 3. Comparison of the operative data of the patients according to the ALBI score greater and lesser than -2.44.

	ALBI score < -2.44 (n=90) N (%) or mean±standard deviation	ALBI score > -2.44 (n=49) N (%) or mean±standard deviation	P-value
CC time	90.59±36.180	96.04±36.487	0.371
CPB time	134.28±48.12	147.94±65.219	0.406
Re-exploration for bleeding	17 (18.9%)	5 (10.2%)	0.180

ALBI=Albumin-Bilirubin; CC=cross-clamping; CPB=cardiopulmonary bypass

Table 4. Univariable and multivariable analyses of mortality.

	Univariable analyses			Multivariable analyses		
	OR	95% CI	P-value	OR	95% CI	P-value
ALBI score	3.83	1.55 – 9.49	0.004*	3.37	1.09 – 10.5	0.036*
EuroSCORE II	1.25	1.09 – 1.44	0.002*	1.09	0.92 – 1.29	0.331
Age (years)	1.05	0.99 – 1.1	0.077			
NYHA class 3-4	4.51	1.38 – 14.86	0.013*	3.14	0.71 – 14.5	0.134
LVEF	0.98	0.93 – 1.04	0.542			
PAP	1.02	0.99 – 1.05	0.187			
DM	1.18	0.35 – 3.97	0.783			
Renal failure	2.83	0.87 – 9.17	0.082			
CPB time	1.02	1.01 – 1.03	< 0.001*	1.02	1.01 – 1.03	0.001*
Combined aortic, mitral, and tricuspid valve surgery	6.78	2.04 – 22.5	0.002*	1.23	0.31 – 6.99	0.851
Infective endocarditis surgery	2.82	0.79 – 10.1	0.109			
Reoperation	2.20	0.75 – 3.92	0.115			

*P-value < 0.05 is statistically significant

ALBI=Albumin-Bilirubin; CI=confidence interval; CPB=cardiopulmonary bypass; DM=diabetes mellitus; EuroSCORE=European System for Cardiac Operative Risk Evaluation; LVEF=left ventricular ejection fraction; NYHA=New York Heart Association; OR=odds ratio; PAP=pulmonary artery pressure

Table 5. Comparison of postoperative complications according to the ALBI score greater and lesser than -2.44.

	ALBI score < -2.44 (n=90) N (%) or mean±standard deviation	ALBI score > -2.44 (n=49) N (%) or mean±standard deviation	P-value
Hospital mortality	4 (4.4%)	12 (24.5%)	< 0.001*
Ventilation time (days)	2.23±4.53	5.16±8.27	0.214
ICU stay (days)	3.89±7.93	7.31±11.46	0.234
Hospital stay (days)	11.83±9.75	15.14±16.38	0.667
Temporary pacemaker requirement	14 (15.6%)	7 (14.2%)	0.842
Permanent pacemaker requirement	8 (8.8%)	3 (6.1%)	0.564
AKI	19 (21.1%)	19 (38.8%)	0.028*
Wound complication	10 (11.1%)	6 (12.2%)	0.841
ECMO	0 (0%)	2 (4.1%)	0.054
IABP	3 (3.3%)	2 (4.1%)	0.821
Postoperative AF	16 (17.8%)	9 (18.4%)	0.931

*P-value < 0.05 is statistically significant.

AF=atrial fibrillation; AKI=acute kidney injury; ALBI=Albumin-Bilirubin; ECMO=extracorporeal membrane oxygenation; IABP=intra-aortic balloon pump; ICU=intensive care unit

In our study, we found that the optimal ALBI cutoff value was -2.44, analyzed by the ROC curve to hospital mortality. This value is very close to the cutoff value (-2.6) between grades 1 and 2 in hepatectomy patients. We found that patients who were operated for infective endocarditis had NYHA classes 3 and 4 complaints, and those who had high PAP had higher ALBI scores. Low albumin

levels, a negative acute phase reactant, due to sepsis developing in infective endocarditis patients cause an increase in ALBI score^[13]. As expected, in our study, ALBI scores were higher in infective endocarditis cases.

We found a positive correlation between ALBI score and PAP. Every 1 change in ALBI score causes a 7.95 mmHg change in PAP. The

relationship between high PAP and mortality in valve surgery has been shown in studies^[14]. The correlation between PAP and ALBI score shows that ALBI score can be used safely in valve surgery to predict mortality as well. Also, as it is known, ALBI score has been shown to reflect right heart failure and related liver dysfunction in patients with acute heart failure^[15]. Therefore, we suggest the use of ALBI score preoperatively to assess surgical risk.

Our study is the first to demonstrate the use of the ALBI scoring system in heart valve surgery to our knowledge. In our study, it was observed that patients with high ALBI scores in multivariate analysis had a higher mortality rate. In multivariate analysis, the ALBI score is a stronger predictor of hospital mortality compared to EuroSCORE II. Also, we think that the ALBI score is clinically important in the preoperative risk assessment phase in heart valve surgery because since more than one third of our study population has a high ALBI score, it is a strong variable in predicting postoperative hospital mortality and is easily measurable.

ALBI is a scoring system that only includes albumin and bilirubin levels of a patient. It's easy to measure and calculate. Therefore, it is valuable for cardiac valvular surgical patients to foresee and predict mortality. Both albumin and bilirubin values are reachable and easy measures to calculate ALBI score. Previous scoring systems such as MELD are poor predictors of outcomes for cardiac patients^[2]. Therefore, ALBI is an easy and cost-effective way to predict the mortality and can be used confidently in valvular surgery patients.

Limitations

Our study has some limitations. First, it only included data from a single-center and a small patient population. Due to the small population, hospital mortality and morbidity were relatively low. We did not investigate the long-term consequences of ALBI score. There is a need for multicenter and prospective studies to support the use of ALBI score, which may be a cost-effective way to predict mortality.

CONCLUSION

ALBI is a strong variable in predicting hospital mortality and AKI in cardiac valvular surgery. It is easy to measure both albumin and bilirubin to calculate ALBI score. We recommend the use of ALBI score preoperatively to predict mortality.

ACKNOWLEDGEMENTS

Both authors were at Mehmet Akif Ersoy Thoracic and Cardiovascular Training and Research Hospital at the time the study was conducted.

And both authors want to thank Dr. Murat Sargin for his support and contribution for the study.

No financial support:

No conflict of interest.

Authors' Roles & Responsibilities

ZMD	Substantial contributions to the design of the work; and the analysis of data for the work; drafting the work; final approval of the version to be published
BT	Substantial contributions to the design of the work; revising the work critically for important intellectual content; final approval of the version to be published

REFERENCES

- Møller S, Bernardi M. Interactions of the heart and the liver. *Eur Heart J*. 2013;34(36):2804-11. doi:10.1093/eurheartj/ehd246.
- Assenza GE, Graham DA, Landzberg MJ, Valente AM, Singh MN, Bashir A, et al. MELD-XI score and cardiac mortality or transplantation in patients after fontan surgery. *Heart*. 2013;99(7):491-6. doi:10.1136/heartjnl-2012-303347.
- Ailawadi G, Lapar DJ, Swenson BR, Siefert SA, Lau C, Kern JA, et al. Model for end-stage liver disease predicts mortality for tricuspid valve surgery. *Ann Thorac Surg*. 2009;87(5):1460-7; discussion 1467-8. doi:10.1016/j.athoracsur.2009.01.043.
- Matthews JC, Pagani FD, Haft JW, Koelling TM, Naftel DC, Aaronson KD. Model for end-stage liver disease score predicts left ventricular assist device operative transfusion requirements, morbidity, and mortality. *Circulation*. 2010;121(2):214-20. doi:10.1161/CIRCULATIONAHA.108.838656.
- Johnson PJ, Berhane S, Kagebayashi C, Satomura S, Teng M, Reeves HL, et al. Assessment of liver function in patients with hepatocellular carcinoma: a new evidence-based approach—the ALBI grade. *J Clin Oncol*. 2015;33(6):550-8. doi:10.1200/JCO.2014.57.9151.
- Han S, Wang C, Tong F, Li Y, Li Z, Sun Z, et al. Prognostic impact of albumin-bilirubin score on the prediction of in-hospital mortality in patients with heart failure: a retrospective cohort study. *BMJ Open*. 2022;12(1):e049325. doi:10.1136/bmjopen-2021-049325.
- Yamada S, Kaneshiro T, Yoshihisa A, Nodera M, Amami K, Nehashi T, et al. Albumin-bilirubin score for prediction of outcomes in heart failure patients treated with cardiac resynchronization therapy. *J Clin Med*. 2021;10(22):5378. doi:10.3390/jcm10225378.
- The Society of Thoracic Surgeons. STS Adult Cardiac Surgery Database Data Specifications. Version 4.20.2. Chicago (IL): The Society of Thoracic Surgeons; c2022. Available at: https://www.sts.org/sites/default/files/ACSD_DataSpecifications_V4_20_2.pdf
- Laribi S, Mebazaa A. Cardiohepatic syndrome: liver injury in decompensated heart failure. *Curr Heart Fail Rep*. 2014;11(3):236-40. doi:10.1007/s11897-014-0206-8.
- Hawkins RB, Young BAC, Mehaffey JH, Speir AM, Quader MA, Rich JB, et al. Model for end-stage liver disease score independently predicts mortality in cardiac surgery. *Ann Thorac Surg*. 2019;107(6):1713-9. doi:10.1016/j.athoracsur.2018.12.011.
- Ma T, Li QS, Wang Y, Wang B, Wu Z, Lv Y, et al. Value of pretransplant albumin-bilirubin score in predicting outcomes after liver transplantation. *World J Gastroenterol*. 2019;25(15):1879-89. doi:10.3748/wjg.v25.i15.1879.
- Zhang ZQ, Xiong L, Zhou JJ, Miao XY, Li QL, Wen Y, et al. Ability of the ALBI grade to predict posthepatectomy liver failure and long-term survival after liver resection for different BCLC stages of HCC. *World J Surg Oncol*. 2018;16(1):208. doi:10.1186/s12957-018-1500-9.
- Arques S. Human serum albumin in cardiovascular diseases. *Eur J Intern Med*. 2018;52:8-12. doi:10.1016/j.ejim.2018.04.014.
- Schubert SA, Mehaffey JH, Booth A, Yarbboro LT, Kern JA, Kennedy JLW, et al. Pulmonary-systemic pressure ratio correlates with morbidity in cardiac valve surgery. *J Cardiothorac Vasc Anesth*. 2019;33(3):677-82. doi:10.1053/j.jvca.2018.08.190.
- Matsue Y, Kagiya N, Yamaguchi T, Kuroda S, Okumura T, Kida K, et al. Clinical and prognostic values of ALBI score in patients with acute heart failure. *Heart Lung Circ*. 2020;29(9):1328-37. doi:10.1016/j.hlc.2019.12.003.

ERRATUM

In the article "Albumin-Bilirubin Score: A Novel Mortality Predictor in Valvular Surgery", DOI <https://doi.org/10.21470/1678-9741-2022-0008>, published in the Brazilian Journal of Cardiovascular Surgery, 38(2):271-277, the references section was incorrectly presented.

In the published article, the references were:

- Hyde JA, Chinn JA, Graham TR. Platelets and cardiopulmonary bypass. *Perfusion*. 1998;13(6):389-407. doi:10.1177/026765919801300603.
- Bønding Andreasen J, Hvas AM, Ravn HB. Marked changes in platelet count and function following pediatric congenital heart surgery. *Paediatr Anaesth*. 2014;24(4):386-92. doi:10.1111/pan.12347.
- Ignjatovic V, Than J, Summerhayes R, Newall F, Horton S, Cochrane A, et al. The quantitative and qualitative responses of platelets in pediatric patients undergoing cardiopulmonary bypass surgery. *Pediatr Cardiol*. 2012;33(1):55-9. doi:10.1007/s00246-011-0079-5.
- Kunitomo R, Tsurusaki S, Suzuki R, Takaji K, Moriyama S, Hagio K, et al. Predictive factors for platelet number after cardiopulmonary bypass and postoperative blood loss. *ASAIO J*. 2002;48(6):671-4. doi:10.1097/00002480-200211000-00018.
- Kertai MD, Zhou S, Karhausen JA, Cooter M, Jooste E, Li YJ, et al. Platelet counts, acute kidney injury, and mortality after coronary artery bypass grafting surgery. *Anesthesiology*. 2016;124(2):339-52. doi:10.1097/ALN.0000000000000959.
- Griffin BR, Bronsart M, Reece TB, Pal JD, Cleveland JC, Fullerton DA, et al. Thrombocytopenia after cardiopulmonary bypass is associated with increased morbidity and mortality. *Ann Thorac Surg*. 2020;110(1):50-7. doi:10.1016/j.athoracsur.2019.10.039.
- Karhausen JA, Smeltz AM, Akushevich I, Cooter M, Podgoreanu MV, Stafford-Smith M, et al. Platelet counts and postoperative stroke after coronary artery bypass grafting surgery. *Anesth Analg*. 2017;125(4):1129-39. doi:10.1213/ANE.0000000000002187.
- Dewar DC, Tarrant SM, King KL, Balogh ZJ. Changes in the epidemiology and prediction of multiple-organ failure after injury. *J Trauma Acute Care Surg*. 2013;74(3):774-9. doi:10.1097/TA.0b013e31827a6e69.
- Gao S, Li Y, Diao X, Yan S, Liu G, Liu M, et al. Vacuum-assisted venous drainage in adult cardiac surgery: a propensity-matched study. *Interact Cardiovasc Thorac Surg*. 2020;30(2):236-42. doi:10.1093/icvts/ivz253.
- Spiezia L, Di Gregorio G, Campello E, Maggiolo S, Bortolussi G, Stellin G, et al. Predictors of postoperative bleeding in children undergoing cardiopulmonary bypass: a preliminary Italian study. *Thromb Res*. 2017;153:85-9. doi:10.1016/j.thromres.2017.03.021.
- Stravitz RT, Ellerbe C, Durkalski V, Reuben A, Lisman T, Lee WM, et al. Thrombocytopenia is associated with multi-organ system failure in patients with acute liver failure. *Clin Gastroenterol Hepatol*. 2016;14(4):613-20.e4. doi:10.1016/j.cgh.2015.09.029.
- Nydam TL, Kashuk JL, Moore EE, Johnson JL, Burlew CC, Biffi WL, et al. Refractory postinjury thrombocytopenia is associated with multiple organ failure and adverse outcomes. *J Trauma*. 2011;70(2):401-6. doi:10.1097/TA.0b013e31820b5c85.
- Greco E, Lupia E, Bosco O, Vizio B, Montrucchio G. Platelets and multi-organ failure in sepsis. *Int J Mol Sci*. 2017;18(10):2200. doi:10.3390/ijms18102200.
- Jansen MPB, Florquin S, Roelofs JTH. The role of platelets in acute kidney injury. *Nat Rev Nephrol*. 2018;14(7):457-71. doi:10.1038/s41581-018-0015-5.
- Bdeir K, Gollomp K, Stasiak M, Mei J, Papiewska-Pajak I, Zhao G, et al. Platelet-specific chemokines contribute to the pathogenesis of acute lung injury. *Am J Respir Cell Mol Biol*. 2017;56(2):261-70. doi:10.1165/rncmb.2015-0245OC.
- Spieess BD. Platelets reflect biologic complexity. *Anesthesiology*. 2016;124(2):265-6. doi:10.1097/ALN.0000000000000960.
- Karkouti K, Wijeyesundera DN, Yau TM, Beattie WS, Abdelnaem E, McCluskey SA, et al. The independent association of massive blood loss with mortality in cardiac surgery. *Transfusion*. 2004;44(10):1453-62. doi:10.1111/j.1537-2995.2004.04144.x.
- Karkouti K. Transfusion and risk of acute kidney injury in cardiac surgery. *Br J Anaesth*. 2012;109 Suppl 1:i29-i38. doi:10.1093/bja/aes422.
- Leytin V, Allen DJ, Mykhaylov S, Mis L, Lyubimov EV, Garvey B, et al. Pathologic high shear stress induces apoptosis events in human platelets. *Biochem Biophys Res Commun*. 2004;320(2):303-10. doi:10.1016/j.bbrc.2004.05.166.
- Tran PL, Pietropaolo MG, Valerio L, Brengle W, Wong RK, Kazui T, et al. Hemolysate-mediated platelet aggregation: an additional risk mechanism contributing to thrombosis of continuous flow ventricular assist devices. *Perfusion*. 2016;31(5):401-8. doi:10.1177/0267659115615206.
- Mohebbi D, Kaplan D, Carlisle M, Supiano MA, Rondina MT. Alterations in platelet function during aging: clinical correlations with thromboinflammatory disease in older adults. *J Am Geriatr Soc*. 2014;62(3):529-35. doi:10.1111/jgs.12700.
- Balduini CL, Noris P. Platelet count and aging. *Haematologica*. 2014;99(6):953-5. doi:10.3324/haematol.2014.106260.
- Leguyader A, Watanabe R, Berbé J, Boumediene A, Cogné M, Laskar M. Platelet activation after aortic prosthetic valve surgery. *Interact Cardiovasc Thorac Surg*. 2006;5(1):60-4. doi:10.1510/icvts.2005.115733.
- Mahmood S, Bilal H, Zaman M, Tang A. Is a fully heparin-bonded cardiopulmonary bypass circuit superior to a standard cardiopulmonary bypass circuit? *Interact Cardiovasc Thorac Surg*. 2012;14(4):406-14. doi:10.1093/icvts/ivr124.
- Jacobson J. Nitric oxide: platelet protectant properties during cardiopulmonary bypass/ECMO. *J Extra Corpor Technol*. 2002;34(2):144-7.
- Selleng S, Selleng K. Heparin-induced thrombocytopenia in cardiac surgery and critically ill patients. *Thromb Haemost*. 2016;116(5):843-51. doi:10.1160/TH16-03-0230.

The correct references are:

- Møller S, Bernardi M. Interactions of the heart and the liver. *Eur Heart J*. 2013;34(36):2804-11. doi:10.1093/eurheartj/ehd246.
- Assenza GE, Graham DA, Landzberg MJ, Valente AM, Singh MN, Bashir A, et al. MELD-XI score and cardiac mortality or transplantation in patients after fontan surgery. *Heart*. 2013;99(7):491-6. doi:10.1136/heartjnl-2012-303347.
- Ailawadi G, Lapar DJ, Swenson BR, Siefert SA, Lau C, Kern JA, et al. Model for end-stage liver disease predicts mortality for tricuspid valve surgery. *Ann Thorac Surg*. 2009;87(5):1460-7; discussion 1467-8. doi:10.1016/j.athoracsur.2009.01.043.
- Matthews JC, Pagani FD, Haft JW, Koelling TM, Naftel DC, Aaronson KD. Model for end-stage liver disease score predicts left ventricular assist device operative transfusion requirements, morbidity, and mortality. *Circulation*. 2010;121(2):214-20. doi:10.1161/CIRCULATIONAHA.108.838656.
- Johnson PJ, Berhane S, Kagebayashi C, Satomura S, Teng M, Reeves HL, et al. Assessment of liver function in patients with hepatocellular carcinoma: a new evidence-based approach—the ALBI grade. *J Clin Oncol*. 2015;33(6):550-8. doi:10.1200/JCO.2014.57.9151.
- Han S, Wang C, Tong F, Li Y, Li Z, Sun Z, et al. Prognostic impact of albumin-bilirubin score on the prediction of in-hospital mortality in patients with heart failure: a retrospective cohort study. *BMJ Open*. 2022;12(1):e049325. doi:10.1136/bmjopen-2021-049325.
- Yamada S, Kaneshiro T, Yoshihisa A, Nodera M, Amami K, Nehashi T, et al. Albumin-bilirubin score for prediction of outcomes in heart failure patients treated with cardiac resynchronization therapy. *J Clin Med*. 2021;10(22):5378. doi:10.3390/jcm10225378.
- The Society of Thoracic Surgeons. STS Adult Cardiac Surgery Database Data Specifications. Version 4.20.2. Chicago (IL): The Society of Thoracic Surgeons; c2022. Available at: https://www.sts.org/sites/default/files/ACSD_DataSpecifications_V4_20_2.pdf
- Laribi S, Mebazaa A. Cardiohepatic syndrome: liver injury in decompensated heart failure. *Curr Heart Fail Rep*. 2014;11(3):236-40. doi:10.1007/s11897-014-0206-8.
- Hawkins RB, Young BAC, Mehaffey JH, Speir AM, Quader MA, Rich JB, et al. Model for end-stage liver disease score independently predicts mortality in cardiac surgery. *Ann Thorac Surg*. 2019;107(6):1713-9. doi:10.1016/j.athoracsur.2018.12.011.
- Ma T, Li QS, Wang Y, Wang B, Wu Z, Lv Y, et al. Value of pretransplant albumin-bilirubin score in predicting outcomes after liver transplantation. *World J Gastroenterol*. 2019;25(15):1879-89. doi:10.3748/wjg.v25.i15.1879.
- Zhang ZQ, Xiong L, Zhou JJ, Miao XY, Li QL, Wen Y, et al. Ability of the ALBI grade to predict posthepatectomy liver failure and long-term survival after liver resection for different BCLC stages of HCC. *World J Surg Oncol*. 2018;16(1):208. doi:10.1186/s12957-018-1500-9.
- Arques S. Human serum albumin in cardiovascular diseases. *Eur J Intern Med*. 2018;52:8-12. doi:10.1016/j.ejim.2018.04.014.
- Schubert SA, Mehaffey JH, Booth A, Yarburo LT, Kern JA, Kennedy JLW, et al. Pulmonary-systemic pressure ratio correlates with morbidity in cardiac valve surgery. *J Cardiothorac Vasc Anesth*. 2019;33(3):677-82. doi:10.1053/j.jvca.2018.08.190.
- Matsue Y, Kagiya N, Yamaguchi T, Kuroda S, Okumura T, Kida K, et al. Clinical and prognostic values of ALBI score in patients with acute heart failure. *Heart Lung Circ*. 2020;29(9):1328-37. doi:10.1016/j.hlc.2019.12.003.

The Brazilian Journal of Cardiovascular Surgery apologizes for this error, which has now been corrected.